

Paleoenvironmental conditions in the Spanish Miocene–Pliocene boundary: isotopic analyses of *Hipparion* dental enamel

Laura Domingo · Stephen T. Grimes ·
M. Soledad Domingo · M. Teresa Alberdi

Received: 28 May 2008 / Revised: 5 December 2008 / Accepted: 5 December 2008
© Springer-Verlag 2009

Abstract Expansion of C₄ grasses during Late Miocene and Early Pliocene constitutes one of the most remarkable biotic events of the Cenozoic era. The Teruel–Alfambra region (northeastern Spain) contains one of the most complete Miocene–Pliocene sequences of mammalian fossil sites in the world. In this study, stable isotope ($\delta^{13}\text{C}$ and $\delta^{18}\text{O}$) analyses have been performed on the tooth enamel from the equid *Hipparion* from 19 localities spanning a time interval from approximately 10.9 to 2.7 Ma. This time range starts with the first appearance of this genus in Spain and ends at its extinction. An increase in $\delta^{13}\text{C}$ at about 4.2 Ma has been observed, indicative of a shift toward a more open habitat. This shift may be related to a large scale vegetation change which occurred across the Miocene–Pliocene boundary when C₄ grasses expanded. This expansion might in turn be linked to global tectonic events such as the uplift of the Himalaya and/or the closure

of the Panama Isthmus. However, other more regional factors may have ultimately enhanced the trend toward more open habitats in the Western Mediterranean Basin. The Messinian Salinity Crisis was a major environmental event that may have been responsible for the isotopic changes seen in the equid *Hipparion* from the Iberian Peninsula along with an increase in the aridity detected ~4.6 Ma ago in the Sahara. Even though the exact factor triggering the isotopic change observed in the *Hipparion* enamel remains mostly unknown, this study demonstrates that the global environmental changes detected across the Miocene–Pliocene boundary are also recorded in the realm of the Iberian Peninsula.

Keywords *Hipparion* · Carbon and oxygen stable isotopes · Miocene–Pliocene boundary · Teruel–Alfambra region

Electronic supplementary material The online version of this article (doi:10.1007/s00114-008-0500-y) contains supplementary material, which is available to authorized users.

L. Domingo (✉)
Departamento de Paleontología, Facultad CC. Geológicas,
Universidad Complutense de Madrid,
28040 Madrid, Spain
e-mail: lauradomingo@geo.ucm.es

S. T. Grimes
School of Earth, Ocean and Environmental Sciences,
University of Plymouth,
Drake Circus,
Plymouth, Devon PL4 8AA, UK

M. S. Domingo · M. T. Alberdi
Departamento de Paleobiología,
Museo Nacional de Ciencias Naturales (CSIC),
28006 Madrid, Spain

Introduction

The carbon isotope composition of enamel from mammal herbivores has been used to distinguish between different types of paleodiets as it faithfully records the isotopic composition of consumed plants (DeNiro and Epstein 1978; Lee-Thorp and van der Merwe 1987; Quade et al. 1992), which depends on the photosynthetic pathways they use. According to this, plants can be grouped into three different photosynthetic pathways: C₃, C₄, and crassulacean acid metabolism (CAM). C₃ plants (85% of all terrestrial plants) use the Calvin photosynthetic pathway and include trees, shrubs, forbs, and cool-season grasses. C₄ plants (5–10% of all terrestrial plants) use the Hatch–Slack photosynthetic pathway and are mainly comprised of tropical, warm-season sedges, and grasses. Finally, the CAM photosynthetic pathway is used by a group of plants adapted to dry

conditions. C_3 plants discriminate more against the heavy isotope (^{13}C) during photosynthetic fixation of CO_2 compared to C_4 plants. This results in very different carbon isotopic compositions between these two kinds of photosynthetic pathways. $\delta^{13}\text{C}$ values of C_3 plants range from -34‰ to -23‰ (VPDB), with an average of -27‰ , whereas C_4 plants have $\delta^{13}\text{C}$ values ranging from -17‰ to -9‰ (VPDB), with an average of -13‰ (Bender 1971; Farquhar et al. 1989). CAM plants show intermediate carbon isotopic values. Nevertheless, this group of plants is not very frequently taken into account in paleodietary studies as they have rarely constituted an important food source for herbivore mammals.

Cerling and Harris (1999) proposed an enrichment in ^{13}C between tooth enamel and diet of $\sim 14\text{‰}$ in such a way that tooth enamel from herbivores feeding on C_3 vegetation have $\delta^{13}\text{C}$ values between -16‰ and -9‰ , whereas those eating C_4 grasses display $\delta^{13}\text{C}$ values between -1‰ and $+3\text{‰}$.

Oxygen isotopes from enamel apatite have been used to obtain information regarding paleotemperatures and the evolution of past climates (Ayliffe et al. 1992; Sánchez Chillón et al. 1994; Lécuyer et al. 2003). The $\delta^{18}\text{O}$ value in both enamel phosphate and carbonate is related to that of body water, which in turn record water uptake (inspired O_2 , drinking water, and plant water) and loss (excretion, expired CO_2 , and water vapor) during tooth development (Bryant and Froelich 1995). By analyzing $\delta^{18}\text{O}$ values from herbivore tooth enamel, it is possible to differentiate between obligate drinkers (animals that obtain most of their water from drinking, frequently grazers) and non-obligate drinkers (those obtaining water mainly from plants, frequently browsers; Kohn 1996; Levin et al. 2006). However, when analyzing only one genus (as occurs in this study), enamel $\delta^{18}\text{O}$ can be used to track changes in the isotopic composition of meteoric water, which in turn is positively correlated to mean annual temperature (Dansgaard 1964; Rozanski et al. 1993). In this way, an increase in temperature and aridity is associated with an enrichment in the heavy isotope (^{18}O), whereas a decrease in temperature and greater humidity results in a depletion of the heavy isotope (Koch et al. 1989; Ayliffe et al. 1992).

A significant shift in global terrestrial environments occurred during late Miocene and early Pliocene epochs (between ~ 7 to ~ 5 Ma) characterized by the expansion of C_4 grasses. The predominance of the C_3 versus C_4 plants depends on temperature and CO_2 partial pressure (Ehleringer and Pearcy 1983; Ehleringer and Monson 1993). Ecosystems dominated by C_4 plants are those which display high temperatures during the growing season and aridity. They are also those which display low CO_2 partial pressure values as C_4 plants show anatomical and biochemical adaptations that allow them to concentrate the

CO_2 in the leaf cells before carrying out photosynthesis (Ehleringer et al. 1997). On account of the high temperatures, C_4 -dominated ecosystems appeared first in low latitudes before extending to higher latitudes. The change from C_3 dominated ecosystems to those dominated by C_4 plants was initially studied in paleosol carbonates from the sediments of the Siwaliks (Pakistan) (Quade et al. 1992; Morgan et al. 1994). A sudden change in $\delta^{13}\text{C}$ was observed between 7 and 5 Ma. Subsequently, several studies have demonstrated that the expansion of C_4 plants can also be detected in other sections from Africa, Asia, North America, and South America (e.g., Quade et al. 1992; Morgan et al. 1994; Cerling et al. 1997; Latorre et al. 1997; Fox and Koch 2004; Wang et al. 2006). This expansion was initially proposed to have been a consequence of a global reduction in the atmospheric CO_2 concentration in the late Miocene (Cerling et al. 1993). Other authors have suggested that more gradual regional controls may have also played an important role on account of the lack of synchronicity in the global expansion of C_4 plants (Morgan et al. 1994; Pagani et al. 1999; Tiplle and Pagani 2007).

Europe has been poorly represented in the global study of C_4 plant expansion as there are very few sections with a complete mammalian record spanning the Miocene–Pliocene boundary (Cerling et al. 1997). Nevertheless, the Teruel–Alfambra region in northeastern Spain (Fig. 1) constitutes an excellent location to study the vegetation shift that took place during the late Neogene as it contains thousands of remains of vertebrates from different paleontological sites with well-constrained ages. In this paper, we show enamel $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values from the equid *Hipparion* from 19 paleontological sites spanning an age range between approximately 10.9 and 2.7 Ma.


Geological setting, samples, and methods

The Teruel–Alfambra region is located in the northern part of the Teruel Basin. This region belongs to the middle mountainous Mediterranean climatic domain. In the present day, the vegetation is fully dominated by C_3 plants, although a minor C_4 component (e.g., *Salsola vermiculata*) can be found in the Teruel–Alfambra region and in the nearby Ebro Basin.

The Teruel Basin is situated in the northeastern side of the Iberian Peninsula (Fig. 1) and is oriented NNE–SSW. The basin is $\sim 100\text{-km}$ long and $\sim 15\text{-km}$ wide and is filled by a complete succession spanning middle to late Neogene sediments which are more than 500 m in thickness (van Dam et al. 2001).

The Neogene succession from the Teruel–Alfambra region is bounded by siliciclastic and evaporitic formations

Fig. 1 Geological setting of the Teruel–Alfambra region (1 in the Iberian Peninsula map, which shows the location of Tertiary basins in gray) with the paleontological sites selected in this study: ROM La Roma 2, MB Masía del Barbo 2B, VIP Vivero de Pinos, PM Puente Minero, ALJ Los Aljezares, MDV Masada del Valle 2, CC2 Las Pedrizas, LM Los Mansuetos, CG Cerro de la Garita, ARQ El Arquillo 1, KS Las Casiones, MIL Milagros, PER Peralejos E, VB Villalba Alta. (Modified from van Dam et al. 2001). The La Calera and Orrios sites do not appear on the map and are situated toward the South and North of the basin, respectively. The Nombrevilla 1 and Villarroya sites belong to the Calatayud and Ebro basins, respectively, meanwhile the Layna site is situated in the Iberian Range (2, 3, and 4, respectively, in the Iberian Peninsula map)



of Triassic age and also by Jurassic aged carbonate deposits. The oldest Neogene deposits are situated in the eastern part of the basin and are Early Miocene in age. In the middle part of the basin (Alfambra River valley), sediments are Late Miocene to Early Pliocene in age and represent alluvial fan distal facies and shallow lacustrine environments (van Dam et al. 2001). The Neogene deposits overlie unconformably upon Mesozoic and Paleogene formations. The dating method used in this area is mainly based on mammal faunas which are widely distributed across the basin (Alcalá 1994). The majority of the mammal localities are found in stratigraphic superposition. Several sections have also been dated by using magnetostratigraphic analysis (Alcalá et al. 2000). This allows correlation between the biostratigraphy of the mammalian succession and the geomagnetic polarity time scale (GPTS) of Cande and Kent (1995).

Tooth enamel samples, 134, from seven different species of the equid *Hipparion* (*Hipparion primigenium*, *Hipparion matthewi*, *Hipparion periafricanum*, *Hipparion laromae*, *Hipparion concudense*, *Hipparion fissurae*, and *Hipparion*

rocinantis) were analyzed for $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$. Samples came from 19 paleontological sites, 16 of which are located in the Teruel–Alfambra region, with the rest belonging to sites in the Calatayud and Ebro basins (Nombrevilla 1 and Villarroya) and the Iberian Range (Layna; Table 1 and Electronic supplementary material S1). Paleogeographic studies demonstrate that the architecture of the Northern Iberian Peninsula from 12 Ma ago to the present day has not changed substantially (Popov et al. 2004). Ages were assigned to the fossil sites (Table 1) according to Domingo et al. (2007), who carried out a biochronological study of the Upper Neogene of the Iberian Peninsula based on the fauna of macromammals using the maximum likelihood appearance event ordination (ML AEO) method (Alroy 2000).

By using a single genus (*Hipparion*), vital effects (stable isotope fractionation due to different metabolisms), which could affect the interpretation of the isotope results, were reduced. Moreover, the analysis of enamel rather than bone should reduce diagenetic issues as enamel is less prone to diagenetic alteration compared to bone (Ayliffe et al. 1994; Lee-Thorp and Sponheimer 2003). Finally, when

Table 1 Biochronological ordination of the paleontological sites from the Teruel-Alfambra region used in this study and arranged by Domingo et al. (2007) using the *Maximum Likelihood Appearance Event Ordination* (ML AEO) method (Alroy, 2000).

Age (Ma)	Number of samples	Fossil sites	MN unit
2.720	4	Villarroya	MN16
3.912	7	Layna	MN15
4.180–3.580*	7	Villalba Alta	MN15
–	1	Perales E	MN14
4.186	8	Orrios 1	MN14
4.186	7	La Calera	MN14
5.689	7	Milagros	MN13
6.080	10	Las Casiones	MN13
6.319	19	El Arquillo 1	MN13
7.011	9	Cerro de la Garita	MN12
7.011	8	Los Mansuetos	MN12
7.011	3	Las Pedrizas	MN12
7.255	4	Masada del Valle 2	MN12
7.408	7	Los Aljezares	MN12
7.834	7	Puente Minero	MN11
8.040	6	Vivero de Pinos	MN11
8.789	7	Masia del Barbo 2B	MN10
8.789	5	La Roma 2	MN10
10.873	8	Nombrevilla 1	MN9

An absolute age could not be assigned to Villalba Alta (marked with an asterisk) by using this technique. Therefore, the paleomagnetic age proposed by Opdyke et al. (1997) and Oms et al. (1999) has been used. In the same way, an absolute age could not be attributed to Perales E where no paleomagnetic studies have been performed so far. Therefore, it has been placed in the chronological order as proposed by Domingo et al. (2007). Nombrevilla 1 and Villarroya belong to Calatayud and Ebro basins, respectively, and Layna corresponds to a karstic filling situated in the Iberian Range.

possible, third molars were used in this study to overcome paleoecological bias from juvenile development (Bryant et al. 1996).

The enamel was recovered by using a rotary drill with a diamond-tipped dental burr, and in some cases, by hand picking enamel under a binocular microscope from lightly crushed fragments. Enamel was removed from as large an area of the tooth as possible to avoid biasing the isotopic results with respect to seasonality, thus producing a homogeneous signal from several months or years. The material analyzed in this study is archived at Fundación Conjunto Paleontológico de Teruel (Dinópolis, Teruel, Spain), Museo Nacional de Ciencias Naturales (CSIC, Madrid, Spain), and Museo Geominero (Instituto Geológico y Minero de España, Madrid, Spain).

The carbon and oxygen isotopic results are reported in the δ -notation against VPDB. $\delta_{\text{sample}} = [(R_{\text{sample}} - R_{\text{standard}})/R_{\text{standard}}] \times 1,000$; with $R = {}^{13}\text{C}/{}^{12}\text{C}$ and ${}^{18}\text{O}/{}^{16}\text{O}$. The analyses were conducted at the University of Plymouth (United Kingdom) using an acid digestion technique with a continuous helium

flow GV IsoPrime with a multiflow preparation system. Approximately 4–5 mg of raw sample was weighed into individual, septum-sealed vials and placed in a hot plate maintained at 90°C. Each vial was automatically flushed with helium before excess H_3PO_4 was added. After approximately 24 h of equilibration time, the CO_2 was analyzed by continuous flow mass spectrometry. The in run NBS-19 standard gave a $\delta^{13}\text{C}$ value of $2.04 \pm 0.24\text{‰}$ (VPDB; $n=5$) and a $\delta^{18}\text{O}$ value of $-2.86 \pm 0.07\text{‰}$ (VPDB; $n=5$). The NBS 120c Florida phosphate rock provided a $\delta^{13}\text{C}$ value of $-6.21 \pm 0.54\text{‰}$ (VPDB; $n=5$) and a $\delta^{18}\text{O}$ value of $-2.24 \pm 0.52\text{‰}$ (VPDB; $n=5$). The reason why the samples were not pretreated prior to isotopic analysis was mainly due to the small amount of sample that was available. This omission of this stage should not have greatly affected our results as Koch et al. (1997) showed that the isotopic value of untreated enamel provides an average value among the isotopic values furnished by different types of pretreatment. Moreover, Rare Earth Element analyses carried out on the same samples indicated that no major diagenetical alteration has affected the *Hipparion* tooth enamel (Electronic supplementary material S2).

Results

Figure 2 (and Electronic supplementary material S1) shows $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values measured on the tooth enamel of the equid *Hipparion* from the Teruel–Alfambra region, Calatayud and Ebro basins, and the Iberian Range. $\delta^{13}\text{C}$ values range between -13.3‰ and -8.5‰ (with a mean value of $-11.6 \pm 0.8\text{‰}$). $\delta^{18}\text{O}$ values range between -9.9‰ and -3.2‰ (with a mean value of $-6.4 \pm 1.1\text{‰}$). A similar trend exists between $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values (i.e., toward an increase in the isotopic values). However, the correlation coefficient between these two variables is very low ($r=0.26$) suggesting diagenesis has not brought about the homogenization of the isotopic values.

As far as $\delta^{13}\text{C}$ values are concerned, significant statistical differences by means of ANOVA tests have not been detected between different *Hipparion* species ($F_{6, 126}=1.732$, $p=0.119$). This suggests that $\delta^{13}\text{C}$ values have not been biased on account of vital effects. Significant differences have also not been found when comparing different types of teeth (ANOVA, $F_{2, 123}=0.534$, $p=0.588$); therefore, the isotopic results are not affected by differences in the time of mineralization associated with each tooth.

In general, $\delta^{13}\text{C}$ values are typical for a C_3 -based diet. The range of $\delta^{13}\text{C}$ values calculated for the plants, by using the dental enamel $\delta^{13}\text{C}$ values from the *Hipparion* and an isotopic fractionation of around -14‰ (Cerling and Harris 1999) is -26.2 to -23.4‰ (VPDB), which are the typical $\delta^{13}\text{C}$ values for C_3 plants (-34 to -23‰ VPDB). However,




Fig. 2 Chronological scheme for the Miocene–Pliocene boundary in Spain and $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values obtained in the *Hipparion* dental enamel from the Teruel–Alfambra region, Calatayud and Ebro basins and Iberian Range. The mean annual precipitation curve obtained by

van Dam (2006) by studying micromammal assemblages from Europe is also shown. Error bars are ± 1 SD. *Hipparion* species reconstruction by Israel M. Sánchez

statistical analyses show significant differences in the $\delta^{13}\text{C}$ between different sites (ANOVA, $F_{17, 115}=2.486$, $p=0.002$) and between Mammal Neogene (MN) units (ANOVA, $F_{7, 125}=5.285$, $p<0.001$) indicating that variations do exist in the carbon isotope signal recorded in the *Hipparion* enamel over time. The $\delta^{13}\text{C}$ values show a near-constant trend until the end of MN14 where a change has been pinpointed. In this case, a significant difference in $\delta^{13}\text{C}$ values before and after the MN14/MN15 boundary (Student t test, $t_{134}=3.616$, $p=0.002$) is evident. This positive isotope shift may be indicating, on the one hand, a slight increase in the consumption of C_4 plants by *Hipparion*, or on the other hand, a change in C_3 vegetation toward more open habitats, where the evaporation rate would have been higher after MN14. If C_4 consumption is responsible, then the percentage of C_4 plants consumed by each individual can be calculated using the equation proposed by Koch et al. (2004). Using this approach, the maximum C_4 consumption has been estimated at 27.5%, whereas the minimum is 0%.

Significant differences by means of ANOVA tests are not observed either in $\delta^{18}\text{O}$ values among species and types of teeth used ($F_{6, 125}=0.950$, $p=0.462$; $F_{2, 122}=0.029$, $p=0.972$, respectively) indicating that vital effects and time of tooth mineralization have not affected the isotopic signal. Contrary to what happens with $\delta^{13}\text{C}$ values, no statistically significant differences have been detected when considering $\delta^{18}\text{O}$ values between sites and MNs on the whole (ANOVA, $F_{17, 114}=0.381$, $p=0.987$; $F_{7, 124}=0.456$, $p=0.865$, respectively).

Changes in the $\delta^{18}\text{O}$ values of ingested water ($\Delta\delta^{18}\text{O}_{\text{water}}$) and mean annual temperature (ΔMAT) have been estimated

by using the equations proposed by Iacumin et al. (1996) and Yurtsever and Gat (1981) which relate $\delta^{18}\text{O}_{\text{CO}_2}$ and $\delta^{18}\text{O}_{\text{water}}$ and $\delta^{18}\text{O}_{\text{water}}$ and MAT ($^{\circ}\text{C}$), respectively. An increase of $\sim 0.7\text{‰}$ has been calculated for $\delta^{18}\text{O}_{\text{water}}$ which corresponds to an increase of $\sim 2.1^{\circ}\text{C}$ in the MAT when comparing values from MN9 to MN16.

Discussion

The expansion of C_4 plants was initially attributed to a drop in the atmospheric CO_2 concentration in the Late Miocene (Cerling et al. 1993). However, Pagani et al. (1999) argued that the CO_2 concentration showed an increase from a minimum value at 14 Ma (~ 180 ppmv) up to values between 320 and 250 ppmv at the end of the Miocene (9 Ma). Therefore, if the mechanism that triggered the expansion of C_4 grasses was the decrease in atmospheric CO_2 concentrations, then the expansion should have occurred at the beginning of the Miocene. Regional factors or changes in the climatic conditions that could have also favored the expansion of C_4 plants, such as an increase in aridity and a change in the seasonal precipitation rate (precipitation during the warm season), are suggested as reasons for the lack of synchronicity observed in the expansion of these plants (Pagani et al. 1999; Fox and Koch 2004).

The tectonic evolution of the Himalaya has been traditionally invoked as playing a major role in the expansion of C_4 plants as it triggered profound changes in oceanic and atmospheric circulation resulting in an increase

in the seasonality of the precipitation and in aridity levels in mid and high latitude regions in the North Hemisphere (Pagani et al. 1999; Wang et al. 2006). This event brought about, likewise, an enhancement in the silicate-rock weathering resulting in the drawdown of atmospheric CO₂ (Ehleringer and Monson 1993).

Another global factor that may have driven a major climatic change is the closure of the Panama Isthmus. This gradual event took place between 13 and 2.6 Ma and resulted in a decrease in the mixing of Atlantic and Pacific waters that gave rise to the development of the modern Atlantic thermohaline circulation, an increase in the temperature, an enhanced evaporation rate in the North Atlantic, and an increase in the precipitation rates in Northern Hemisphere high latitudes (Haug and Tiedemann 1998; Haug et al. 2001; Lunt et al. 2008). The intensification of the Gulf Stream as a consequence of the reorganization of the ocean circulation introduced warm and saline waters to northern latitudes at ~4.6 Ma (Haug and Tiedemann 1998; Lear et al. 2003). This has been related to a warm period between 4.7 and 3.1 Ma, known as the Pliocene Warm Period (Haug et al. 2001).

Even though the tectonic evolution of the Himalaya and the closure of the Panama Isthmus constitute key factors in global oceanic and atmospheric conditions, it is necessary to look for additional regional factors related with the environmental shifts reflected by the isotopic analyses of the *Hipparion* dental enamel from the Teruel–Alfambra region. During the time period represented in this study (~10.9 to 2.7 Ma), tectonic and environmental changes of paramount importance were taking place in the Mediterranean area which gave rise to the Messinian Salinity Crisis (Hsü et al. 1973). It is generally accepted that the main cause of this crisis was the tectonic closure of the Mediterranean Basin that brought about a progressive reduction of the marine water circulation coming from the Atlantic and which culminated in the formation of thick evaporite deposits. Around 5.96 and 5.59 Ma, the deposition of the “Lower Evaporites” and “Upper Evaporites” and “Lago Mare” (Lake Sea) sediments took place, respectively (Krijgsman et al. 1999). The continuous increase in the ice-volume in the Antarctic since the Middle Miocene may have also played an important role in this crisis as it resulted in a drop in sea level which might have restricted oceanic circulation toward the Mediterranean Basin (Rouchy and Caruso 2006). This crisis had important climatic consequences that not only affected the marine realm, but also the terrestrial realm (Agustí et al. 2006; Fortelius et al. 2006; van der Made et al. 2006). Van Dam (2006) carried out a reconstruction of the precipitation rates in Europe between 12 and 3 Ma based on micromammal assemblages. He argued that after the relatively dry conditions of the Middle Miocene (16–13 Ma), an increase

in the precipitation rate took place between 13 and 11 Ma (MN7–8), reaching a maximum at about 11–10 Ma (beginning of the Vallesian, MN9). Van Dam (2006) also proposed that a decrease in the precipitation rate between 10 and 5 Ma, whose minimum was at 4 Ma (Ruscinian, MN15), took place and resulted in an increase in aridity (Fig. 2). Although this author proposed that the tectonic evolution of the Himalaya was ultimately responsible for the increase in aridity in the Late Miocene and Early Pliocene in Europe, the Messinian Salinity Crisis may have played an important role, boosting the aridity pattern detected in the Western Mediterranean Basin. Furthermore, Tiedemann et al. (1989, 1994) pointed out that at ~4.6 Ma, an increase in aridity took place in the southern Sahara, a fact that could have also influenced the environmental conditions of the Iberian Peninsula.

The *Hipparion* dental enamel from the Teruel–Alfambra region has yielded $\delta^{13}\text{C}$ values typical of a diet based on C₃ plants. The significant increase detected at the MN14/MN15 boundary may be indicative of a slight increase in the consumption of C₄ plants. Nevertheless, Hernández Fernández et al. (2007) have shown the predominance of forested biomes in the late Neogene of the Iberian Peninsula, which hindered the development of C₄-dominated vegetation. Even though these kinds of plants have probably never been abundant in the Mediterranean region, Quade et al. (1994) argued that C₄ plants might have been represented in low proportions as can be deduced from $\delta^{13}\text{C}$ values obtained by these authors in Rhodes. Moreover, it should be noted that *Salsola vermiculata*, a C₄ plant of the Chenopodiaceae family, occurs nowadays in the Teruel–Alfambra region and in the nearby Ebro Basin. If the increase in $\delta^{13}\text{C}$ values shown by the *Hipparion* of the Teruel–Alfambra region is interpreted as suggesting an increase in the percentage of C₄ plants in the diet of this equid at ~4.2 Ma, then a slight expansion of C₄ plants in this area might have taken place subsequent to other regions of the world. This would support the notion that the expansion of these plants was controlled by regional climatic and environmental factors, ultimately driven by tectonically controlled changes in the global climate (Latorre et al. 1997; Cerling et al. 1997; Fox and Koch 2004). Cerling et al. (1997) proposed an age of between 8 and 6 Ma for the expansion of C₄ plants in Asia, Africa, South America, and North America based on the isotopic analyses of the dental enamel of different species of macromammals. Our study and those by Latorre et al. (1997), Passey et al. (2002), Fox and Koch (2004), and Ségalen et al. (2007) suggest that such expansion took place between 6 and 4 Ma at higher latitudes.

However, as pointed out before, the increase in $\delta^{13}\text{C}$ values may also be related to a change in the C₃ ecosystems toward more arid conditions (Quade et al. 1989). This

increase in the aridity pattern may be in response to the climatic and environmental changes produced after the Messinian Salinity Crisis which affected regional precipitation rates (Tiedemann et al. 1989, 1994; van Dam 2006) (Fig. 2). This explanation agrees well with a rise in the hypsodonty index detected in different species of *Hipparion* in the Teruel–Alfambra region (Electronic supplementary material S3; Pesquero 2003; Pesquero et al. 2006), as an adaptive process toward a more abrasive diet.

As far as $\delta^{18}\text{O}$ values are concerned, the gradual increase detected from MN9 to MN16 may be related to an increase in the MAT of $\sim 2.1^\circ\text{C}$. Fluteau et al. (2003) carried out a reconstruction of the climatic conditions that took place as a consequence of the Messinian Salinity Crisis using the LMDz Atmospheric General Circulation model and detected an increase in the MAT of $\sim 2.5^\circ\text{C}$ in response to the desiccation of the Mediterranean Basin. Montuire et al. (2006) also estimated MAT values based upon fossil rodent assemblages in Europe from MN9 to MN15. Some of the sites considered by these authors are situated in the Teruel–Alfambra region or in its immediate vicinity (Nombrevilla, Alfambra, Los Aljezares, and La Gloria 4). The increase in MAT values obtained by Montuire et al. (2006), $\sim 3.4^\circ\text{C}$, is not far from the one obtained from the *Hipparion* enamel in this study. Preliminary results based on the bioclimatic analysis of rodent faunas from the Iberian Peninsula have also shown a significant increase in temperature from MN13 to MN14 (Gómez Cano et al. unpublished data). This increase in $\delta^{18}\text{O}$ values during the Mio–Pliocene transition has also been detected on a global scale in Asia (Quade et al. 1992), Europe (Quade et al. 1994), North America (Fox and Koch 2004), South America (Latorre et al. 1997), and Africa (Morgan et al. 1994), with the most significant change being observed between 8 and 4 Ma. Most studies usually agree that the increase in $\delta^{18}\text{O}$ values registered in the dental enamel and paleosoil carbonates at this time might be indicative of an increase in the MATs and/or an increase in aridity pattern (as shown by $\delta^{13}\text{C}$ values; Latorre et al. 1997; Tippie and Pagani 2007 and references therein).

In summary, we have performed stable isotope analyses ($\delta^{13}\text{C}$ and $\delta^{18}\text{O}$) on the *Hipparion* dental enamel from the Teruel–Alfambra region spanning a broad time interval (from 10.9 to 2.7 Ma). Up to now, Europe had been poorly sampled with the aim of gaining information about the environmental conditions that took place across the Miocene–Pliocene boundary, and therefore, this new study provides valuable information for future comparative works. Our data show an increase in $\delta^{13}\text{C}$ values at ~ 4.2 Ma that may be related either to a slight increase in the C_4 vegetation content and/or to a shift in the C_3 ecosystems toward more open habitats at this period of time. Although several contemporary global and regional

events that could have triggered the environmental change suggested by our isotopic results have been explored, the ultimate causes that gave rise to it are currently difficult to assess.

Acknowledgments This research was supported by the Spanish Ministerio de Educación y Ciencia (project CGL2006-04646/BTE and FPU predoctoral contracts to L. D. and M. S. D.). The work conducted by S. T. G. was supported by a Natural Environmental Research Council (NERC) New Investigators grant (NE/C507237/1). Authors are grateful to L. Alcalá and E. Espílez (Fundación Conjunto Paleontológico de Teruel, Dinópolis, Teruel), B. Sánchez-Chillón (Museo Nacional de Ciencias Naturales-CSIC, Madrid) and S. Menéndez (Museo Geominero, Instituto Geológico y Minero de España, Madrid) for kindly providing the studied material. Numerous sampled teeth were found in excavations conducted by L. Alcalá with the authorization of the Dirección General de Patrimonio Cultural del Gobierno de Aragón and supported by the FOCONTUR project (Research Group E-62, Gobierno de Aragón). Thanks are given to M. D. Pesquero (Museo Nacional de Ciencias Naturales-CSIC, Madrid) for the identification of some teeth and for providing information concerning some of the localities from Teruel–Alfambra region. N. López-Martínez, A. M. Alonso-Zarza, M. Hernández Fernández (Universidad Complutense de Madrid, Spain), D. L. Fox, and S. D. Matson (University of Minnesota, USA) are acknowledged for their valuable comments which substantially improved the manuscript. We also thank the editor T. Czeschlik and three anonymous reviewers for helpful review comments.

The experiments carried out in this research comply with the current laws of Spain and the United Kingdom.

References

- Agustí J, Garcés M, Krijgsman W (2006) Evidence for African-Iberian exchanges during the Messinian in the Spanish mammalian record. *Palaeogeogr Palaeoclimatol Palaeoecol* 238:5–14. doi:10.1016/j.palaeo.2006.03.013
- Alcalá L (1994) Macromamíferos Neógenos de la Fosa de Alfambra-Teruel. Instituto de Estudios Turolenses y Museo Nacional de Ciencias Naturales, CSIC, Teruel
- Alcalá L, Alonso-Zarza AM, Azanza B, Calvo JP, Cañaveras JC, van Dam JA, Garcés M, Krijgsman W, van der Meulen AJ, Morales J, Peláez-Campomanes P, Pérez-González A, Sánchez Moral S, Sancho R, Sanz Rubio E (2000) El registro sedimentario y faunístico de las cuencas de Calatayud-Daroca y Teruel. *Evolución paleoambiental y paleoclimática durante el Neógeno*. *Rev Soc Geol Esp* 13(2):323–343
- Alroy J (2000) New methods for quantifying macroevolutionary patterns and processes. *Paleobiology* 26:707–733. doi:10.1666/0094-8373(2000)026<0707: NMFQMP>2.0.CO;2
- Ayliffe LK, Lister AM, Chivas AR (1992) The preservation of glacial-interglacial climatic signatures in the oxygen isotopes of elephant skeletal phosphate. *Palaeogeogr Palaeoclimatol Palaeoecol* 99:179–191. doi:10.1016/0031-0182(92)90014-V
- Ayliffe LK, Chivas AR, Leakey MG (1994) The retention of primary oxygen isotope compositions of fossil elephant skeletal phosphate. *Geochim Cosmochim Acta* 58:5291–5298. doi:10.1016/0016-7037(94)90312-3
- Bender MM (1971) Variations in the $^{13}\text{C}/^{12}\text{C}$ ratios of plants in relation to the pathway of photosynthetic carbon dioxide fixation. *Phytochemistry* 10:1239–1245. doi:10.1016/S0031-9422(00)84324-1

- Bryant JD, Froelich PN (1995) A model of oxygen isotope fractionation in body water of large mammals. *Geochim Cosmochim Acta* 59:4523–4537. doi:[10.1016/0016-7037\(95\)00250-4](https://doi.org/10.1016/0016-7037(95)00250-4)
- Bryant JD, Froelich PN, Showers WJ, Genna BJ (1996) A tale of two quarries: biologic and taphonomic signatures in the oxygen isotope composition of tooth enamel phosphate from modern and Miocene equids. *Palaios* 11:397–408
- Cande SC, Kent DV (1995) Revised calibration of the geomagnetic polarity timescale for the Late Cretaceous and Cenozoic. *J Geophys Res* 100:6093–6095
- Cerling TE, Harris JM (1999) Carbon isotope fractionation between diet and bioapatite in ungulate mammals and implications for ecological and paleoecological studies. *Oecologia* 120:347–363. doi:[10.1007/s004420050868](https://doi.org/10.1007/s004420050868)
- Cerling TE, Wang Y, Quade J (1993) Expansion of C₄ ecosystems as an indicator of global ecological change in the late Miocene. *Nature* 361:344–345. doi:[10.1038/361344a0](https://doi.org/10.1038/361344a0)
- Cerling TE, Harris JM, MacFadden BJ, Leakey MG, Quade J, Eisenmann V, Ehleringer JR (1997) Global vegetation change through the Miocene/Pliocene boundary. *Nature* 389:153–158. doi:[10.1038/38229](https://doi.org/10.1038/38229)
- Dansgaard W (1964) Stable isotopes in precipitation. *Tellus* 16:436–468
- DeNiro MJ, Epstein S (1978) Influence of diet on the distribution of carbon isotopes in animals. *Geochim Cosmochim Acta* 42:495–506
- Domingo MS, Alberdi MT, Azanza B (2007) A new quantitative biochronological ordination for the Upper Neogene mammalian localities of Spain. *Palaeogeogr Palaeoclimatol Palaeoecol* 255:361–376. doi:[10.1016/j.palaeo.2007.08.004](https://doi.org/10.1016/j.palaeo.2007.08.004)
- Ehleringer JR, Monson RK (1993) Evolutionary and ecological aspects of photosynthetic pathway variation. *Annu Rev Ecol Syst* 24:411–439
- Ehleringer JR, Pearcy RW (1983) Variation in quantum yield for CO₂ uptake among C₃ and C₄ plants. *Plant Physiol* 73:555–559
- Ehleringer JR, Cerling TE, Helliker BR (1997) C₄ photosynthesis, atmospheric CO₂, and climate. *Oecologia* 112:285–299. doi:[10.1007/s004420050311](https://doi.org/10.1007/s004420050311)
- Farquhar GD, Ehleringer JR, Hubick KT (1989) Carbon isotopic discrimination and photosynthesis. *Ann Rev Plant Physiol Plant Mol Biol* 40:503–537. doi:[10.1146/annurev.pp.40.060189.002443](https://doi.org/10.1146/annurev.pp.40.060189.002443)
- Fluteau F, Suc J-P, Fauquette S (2003) Modelling the climatic consequences of the Messinian Salinity Crisis. *Geophys Res Abstr* 5:11387
- Fortelius M, Eronen J, Liu L, Pushkina D, Tesakov A, Vislobokova I, Zhang Z (2006) Late Miocene and Pliocene large land mammals and climatic changes in Eurasia. *Palaeogeogr Palaeoclimatol Palaeoecol* 238:219–227. doi:[10.1016/j.palaeo.2006.03.042](https://doi.org/10.1016/j.palaeo.2006.03.042)
- Fox DL, Koch PL (2004) Carbon and oxygen isotopic variability in Neogene paleosol carbonates: constraints on the evolution of the C₄-grasslands of the Great Plains, USA. *Palaeogeogr Palaeoclimatol Palaeoecol* 207:305–329. doi:[10.1016/j.palaeo.2003.09.030](https://doi.org/10.1016/j.palaeo.2003.09.030)
- Haug GH, Tiedemann R (1998) Effect of the formation of the Isthmus of Panama on Atlantic Ocean thermohaline circulation. *Nature* 393:673–676. doi:[10.1038/31447](https://doi.org/10.1038/31447)
- Haug GH, Tiedemann R, Zahn R, Ravelo AC (2001) Role of Panama uplift on oceanic freshwater balance. *Geology* 29:207–210. doi:[10.1130/0091-7613\(2001\)029<0207:ROPUOO>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<0207:ROPUOO>2.0.CO;2)
- Hernández Fernández M, Álvarez Sierra MA, Peláez-Campomanes P (2007) Bioclimatic analysis of rodent palaeofaunas reveals severe climatic changes in Southwestern Europe during the Plio-Pleistocene. *Palaeogeogr Palaeoclimatol Palaeoecol* 251:500–526. doi:[10.1016/j.palaeo.2007.04.015](https://doi.org/10.1016/j.palaeo.2007.04.015)
- Hsü KJ, Ryan WBF, Cita MB (1973) Late Miocene dessication of the Mediterranean. *Nature* 242:240–244. doi:[10.1038/242240a0](https://doi.org/10.1038/242240a0)
- Iacumin P, Bocherens H, Mariotti A, Longinelli A (1996) Oxygen isotope analyses of co-existing carbonate and phosphate in biogenic apatite: a way to monitor diagenetic alteration of bone phosphate? *Earth Planet Sci Lett* 142:1–6. doi:[10.1016/0012-821X\(96\)00093-3](https://doi.org/10.1016/0012-821X(96)00093-3)
- Koch PL, Fisher DC, Dettman D (1989) Oxygen isotope variation in the tusks of extinct proboscideans: a measure of season of death and seasonality. *Geology* 17:515–519. doi:[10.1130/0091-7613\(1989\)017<0515:OIVITT>2.3.CO;2](https://doi.org/10.1130/0091-7613(1989)017<0515:OIVITT>2.3.CO;2)
- Koch PL, Tuross N, Fogel ML (1997) The effects of sample treatment and diagenesis on the isotopic integrity of carbonate in biogenic hydroxylapatite. *J Archaeol Sci* 24:417–429. doi:[10.1006/jasc.1996.0126](https://doi.org/10.1006/jasc.1996.0126)
- Koch PL, Diffenbaugh NS, Hoppe KA (2004) The effects of Late Quaternary climate and pCO₂ change on C₄ plant abundance in the South-Central United States. *Palaeogeogr Palaeoclimatol Palaeoecol* 207:331–357. doi:[10.1016/j.palaeo.2003.09.034](https://doi.org/10.1016/j.palaeo.2003.09.034)
- Kohn MJ (1996) Predicting animal $\delta^{18}\text{O}$: accounting for diet and physiological adaptation. *Geochim Cosmochim Acta* 60:4811–4829. doi:[10.1016/S0016-7037\(96\)00240-2](https://doi.org/10.1016/S0016-7037(96)00240-2)
- Krijgsman WJ, Hilgen FJ, Raffi I, Sierro FJ, Wilson DS (1999) Chronology, causes and progression of the Messinian salinity crisis. *Nature* 400:652–655. doi:[10.1038/23231](https://doi.org/10.1038/23231)
- Latorre C, Quade J, McIntosh WC (1997) The expansion of C₄ grasses and global change in the late Miocene: stable isotope evidence from the Americas. *Earth Planet Sci Lett* 146:83–96. doi:[10.1016/S0012-821X\(96\)00231-2](https://doi.org/10.1016/S0012-821X(96)00231-2)
- Lear CH, Rosenthal Y, Wright JD (2003) The closing of a seaway: ocean water masses and global climate change. *Earth Planet Sci Lett* 210:425–436. doi:[10.1016/S0012-821X\(03\)00164-X](https://doi.org/10.1016/S0012-821X(03)00164-X)
- Lécuyer C, Picard S, Garcia J-P, Sheppard SMF, Grandjean P, Dromart G (2003) Thermal evolution of Tethyan surface waters during the Middle–Late Jurassic: evidence from $\delta^{18}\text{O}$ values of marine fish teeth. *Paleoceanography* 18:21(1)–21(16). doi:[10.1029/2002PA000863](https://doi.org/10.1029/2002PA000863)
- Lee-Thorp J, Sponheimer M (2003) Three case studies used to reassess the reliability of fossil bone and enamel isotope signals for paleodietary studies. *J Anthropol Archaeol* 22:208–216. doi:[10.1016/S0278-4165\(03\)00035-7](https://doi.org/10.1016/S0278-4165(03)00035-7)
- Lee-Thorp J, van der Merwe NJ (1987) Carbon isotope analysis of fossil bone apatites. *S Afr J Sci* 83:712–715
- Levin NE, Cerling TE, Passey BH, Harris JM, Ehleringer JR (2006) A stable isotope aridity index for terrestrial environments. *Proc Natl Acad Sci* 103:11201–11205. doi:[10.1073/pnas.0604719103](https://doi.org/10.1073/pnas.0604719103)
- Lunt DJ, Valdes PJ, Haywood A, Rutt IC (2008) Closure of the Panama Seaway during the Pliocene: implications for climate and Northern Hemisphere glaciation. *Clim Dyn* 30:1–18. doi:[10.1007/s00382-007-0265-6](https://doi.org/10.1007/s00382-007-0265-6)
- Montuire S, Maridet O, Legendre S (2006) Late Miocene–Early Pliocene temperature estimates in Europe using rodents. *Palaeogeogr Palaeoclimatol Palaeoecol* 238:247–262. doi:[10.1016/j.palaeo.2006.03.026](https://doi.org/10.1016/j.palaeo.2006.03.026)
- Morgan ME, Kingston JD, Marino BD (1994) Carbon isotopic evidence for the emergence of C₄ plants in the Neogene from Pakistan and Kenya. *Nature* 367:162–165. doi:[10.1038/367162a0](https://doi.org/10.1038/367162a0)
- Oms O, Dinarès-Turell J, Agustí J, Parés JM (1999) Refinements of the European Mammal Biochronology from the magnetic polarity record on the Plio–Pleistocene Zújar Section, Guadix-Baza Basin, SE Spain. *Quat Res* 51:94–103. doi:[10.1006/qres.1998.2018](https://doi.org/10.1006/qres.1998.2018)
- Opdyke N, Mein P, Lindsay A, Pérez-González A, Moissenet E, Norton VL (1997) Continental deposits, magnetostratigraphy

- and vertebrate paleontology, late Neogene of Eastern Spain. *Palaeogeogr Palaeoclimatol Palaeoecol* 133:129–148. doi:[10.1016/S0031-0182\(97\)00080-1](https://doi.org/10.1016/S0031-0182(97)00080-1)
- Pagani M, Arthur MA, Freeman KH (1999) Late Miocene atmospheric CO₂ concentrations and the expansion of the C₄ grass. *Science* 285:876–878. doi:[10.1126/science.285.5429.876](https://doi.org/10.1126/science.285.5429.876)
- Passey BH, Cerling TE, Perkins ME, Voorhies MR, Harris JM, Tucker ST (2002) Environmental change in the Great Plains: an isotopic record from fossil horses. *J Geol* 110:123–140. doi:[10.1086/338280](https://doi.org/10.1086/338280)
- Pesquero MD (2003) *Hipparion* del Turoliense superior de Las Casiones (Fosa de Teruel). *Coloq Paleontol v. ext* 1:511–548
- Pesquero MD, Alberdi MT, Alcalá L (2006) New species of *Hipparion* from La Roma 2 (Late Vallesian; Teruel, Spain): a study of the morphological and biometric variability of *Hipparion primigenium*. *J Paleontol* 80:343–356. doi:[10.1666/0022-3360](https://doi.org/10.1666/0022-3360)
- Popov SV, Rögl F, Rozanov AY, Steininger FF, Shcherba IG, Kováč M (2004) Lithological–paleogeographic maps of Paratethys. 10 maps Late Eocene to Pliocene. *Courier Forschungsinstitut Senckenberg* 250:1–46
- Quade J, Cerling TE, Bowman JR (1989) Systematic variations in the carbon and oxygen isotopic composition of pedogenic carbonate along elevation transects in the southern Great Basin, United States. *Geol Soc Am Bull* 101:464–475
- Quade J, Cerling TE, Barry JC, Morgan ME, Pilbeam DR, Chivas AR, Lee-Thorp JA, van der Merwe NJ (1992) A 16-Ma record of paleodiet using carbon and oxygen isotopes in fossil teeth from Pakistan. *Chem Geol* 94:183–192
- Quade J, Solounias N, Cerling TE (1994) Stable isotopic evidence from paleosol carbonates and fossil teeth in Greece for forest or woodlands over the past 11 Ma. *Palaeogeogr Palaeoclimatol Palaeoecol* 108:41–53. doi:[10.1016/0031-0182\(94\)90021-3](https://doi.org/10.1016/0031-0182(94)90021-3)
- Rouchy JM, Caruso A (2006) The Messinian salinity crisis in the Mediterranean basin: a reassessment of the data and an integrated scenario. *Sediment Geol* 188–189:35–67. doi:[10.1016/j.sedgeo.2006.02.005](https://doi.org/10.1016/j.sedgeo.2006.02.005)
- Rozanski K, Araguás-Araguás L, Gonfiantini R (1993) Isotopic patterns in modern global rainfall. In: Swart PK, Lohmann KC, McKenzie J, Savin S (eds) *Climate Change in continental isotopic records*. *Geophysical Monograph* 78:1–36
- Sánchez Chillón B, Alberdi MT, Leone G, Bonadonna FP, Stenni B, Longinelli A (1994) Oxygen isotopic composition of fossil equid tooth and bone phosphate: an archive of difficult interpretation. *Palaeogeogr Palaeoclimatol Palaeoecol* 107:317–328. doi:[10.1016/0031-0182\(94\)90103-1](https://doi.org/10.1016/0031-0182(94)90103-1)
- Ségalen L, Lee-Thorp JA, Cerling TE (2007) Timing of C₄ grass expansion across sub-Saharan Africa. *J Hum Evol* 53:549–559. doi:[10.1016/j.jhevol.2006.12.010](https://doi.org/10.1016/j.jhevol.2006.12.010)
- Tiedemann R, Sarnthein M, Stein R (1989) Climatic changes in the Western Sahara: aeolo-marine sediment record of the last 8 million years (sites 657–661). *Proc Ocean Drill Prog* 108:241–277
- Tiedemann R, Sarnthein M, Shackleton NJ (1994) Astronomic timescale for the Pliocene Atlantic $\delta^{18}\text{O}$ and dust flux records of Ocean Drilling Program site 659. *Paleoceanography* 9:619–638
- Tipple BJ, Pagani M (2007) The early origins of terrestrial C₄ photosynthesis. *Ann Rev Earth Planet Sci* 35:435–461. doi:[10.1146/annurev.earth.35.031306.140150](https://doi.org/10.1146/annurev.earth.35.031306.140150)
- van Dam JA (2006) Geographic and temporal patterns in the late Neogene (12–3 Ma) aridification of Europe: the use of small mammals as paleoprecipitation proxies. *Palaeogeogr Palaeoclimatol Palaeoecol* 238:190–218. doi:[10.1016/j.palaeo.2006.03.025](https://doi.org/10.1016/j.palaeo.2006.03.025)
- van Dam JA, Alcalá L, Alonso-Zarza AM, Calvo JP, Garcés M, Krijgsman W (2001) The upper Miocene mammal record from the Teruel–Alfambra region (Spain). The MN system and continental stage/age concepts discussed. *J Vertebr Paleontol* 21:367–385
- van der Made J, Morales J, Montoya P (2006) Late Miocene turnover in the Spanish mammal record in relation to palaeoclimate and the Messinian Salinity Crisis. *Palaeogeogr Palaeoclimatol Palaeoecol* 238:228–246. doi:[10.1016/j.palaeo.2006.03.030](https://doi.org/10.1016/j.palaeo.2006.03.030)
- Wang Y, Deng T, Biasatti DM (2006) Ancient diets indicate significant uplift of southern Tibet after ca. 7 Ma. *Geology* 34:309–312. doi:[10.1130/g22254.1](https://doi.org/10.1130/g22254.1)
- Yurtsever Y, Gat J (1981) Atmospheric waters. In: Gat JR, Gonfiantini R (eds) *Stable isotope hydrology. Deuterium and oxygen-18 in the water cycle*. 210. Tech. Rep. Ser. IAEA, Vienna, pp 103–142